



TOOLS 5X EvaGreen qPCR Mix without ROX

Cat. no.: TF-5EVNR250

Storage: –20°C for up to 1 year

Product Size: 1 mL

Introduction

The TOOLS 5× EvaGreen qPCR Mix without ROX is optimized for real-time quantitative polymerase chain reaction (PCR) assays. The ready-to-load mix includes TOOLS HotTaq DNA polymerase, ultrapure dNTPs, MgCl₂, and EvaGreen® dye. Only water, template, and primers need to be added. TOOLS HotTaq DNA polymerase is activated by a 15-min incubation step at 95°C. This prevents extension of nonspecifically annealed primers and primer-dimers formed at low temperatures during qPCR.

EvaGreen® Dye: EvaGreen® is a DNA-binding dye for qPCR that, compared with SYBR® Green I, has similar spectra but much lower PCR inhibition. It is extremely stable and has been shown to be nonmutagenic and noncytotoxic. EvaGreen® is compatible with all common real-time PCR cyclers—simply select the standard settings for SYBR® Green or FAM

EvaGreen® is a registered trademark and licensed for sale by Biotium, Inc.

SYBR® is a registered trademark of Molecular Probes, Inc.

Composition

TOOLS HotTaq DNA polymerase

5× qPCR buffer E

12.5 mM MgCl₂ (1× PCR solution–2.5 mM MgCl₂)

dNTPs, including dTTP to improve reaction sensitivity and efficiency compared with dUTP

EvaGreen® dye

Cycler compatibility

The TOOLS 5× EvaGreen qPCR Mix without ROX can be used with all major qPCR cyclers that do not require a reference dye, including Bio-Rad iQ™ 5, Opticon™, and Opticon™ 2; Chromo 4™, MiniOpticon, CFX96, and CFX384; Corbett Rotor-Gene™ 3000 and Rotor-Gene™ 6000; and Roche LightCycler® 480 and Techne Quantica™.

Recommended qPCR reaction mix

Component	Volume	Concentration
TOOLS 5X EvaGreen qPCR Mix	4 µL	1x
Primer Forward (10 pmol/µl)	0.16–0.5 µL	80–250 nM
Primer Reverse (10 pmol/µl)	0.16–0.5 µL	80–250 nM
DNA template	1–5 µL	1–50 ng/µL
H ₂ O PCR grade	up to 20 µL	
Total	20 µL	

Recommended qPCR cycles

Cycle step	Temp.	Time	Cycles
Initial denaturation	95°C	15 min	1
Denaturation	95°C	15 s	40
Annealing	60°-65°C	20 s	
Elongation	72°C	20 s	

Important: To activate the polymerase, incubate it at 95°C for 15 min at the beginning of the qPCR cycle.

The product is for research only and not for diagnostic and clinical use.